Numbers used in numbering
Collection for publication Nos
used inclusive

625- 699-765- 1024-1025-1026 1027-1099

~~1100-1178 not used~~ 1179-1324 used.

1100-1147-77 1325-45-46 1347-23 24-37-1466-
1495-1540-1707-19-20-32-46 1762-1772-83-1924
1932-47-54 58 2044-63 66-82-89 3013-23-30
3043-44-53-71-90-91 31423 43-44-3265-3324
3345-3423-42 3472-75 34-34 36 3440 41-42

Cocos Isl 225-330-331-412

Spondias purpurea
called tropical plum
Introduced

Psychotria flexuosa Wedd Rubiaceae

Bertiera angustifolia

1339

Adiantum prionophyllum, H.B.K.

= A tetraphyllum, H. B. Willd. Sp. V. 441 (1810). a prionophyllum, H.B.K. Nov. Gen. & Sp. I. 20 (1815); Rob (2), 106

Adiantum riciasum, Presl.

a. Alarconianum, Gaud. Voy.
Bonite Bot. t. 99.

Sida peromicifolia Sam
var humilis (Sav.) K Schum
in Prod Fl Bras XII. 3. 320
(1891) Sida humilis Sav Diss
V 277 t 134 f 2 (1788)

Cocos L nucifera L Sp 1188 Palmae

(○) vide ista

Pilotrichum bijunatum (Sch) Musc
Brid.

1376. Cocos Isl

Mus

Sythopodon rigidus
Hook = Grev.

1412 . Cocos Isl

Rhegogonium spiniforme (L.) Brudh
Musci
Brudh.
1400 Cocos del

Dichroncma Hymenolichen

D sericeum (?)

1423 Cacas Del on the bark of trees

Hypnella (C Müll) Jaeg

H pallescens (Hook) Jaeg

1368 Cocos Ds

Oeloblephanum albidum Bedd.

1477 Cocos dsl

Undetermined mosses from
Cocos Isl

1407

1432

1405-

1431

Eichhornia Crispum

✓247
1470

✓248
1457

✓249
1408-09

Eichhornia L.
E. Crispum L Sp. Pl. 1097 (1753

Hex. St. Ind, Sam. West Africa

Fuchsiones trichoidium Sw. Close
to this species

246
1343 Fairly common on trunks of tree at
300 ft

Fuchsiones L
F Capillaceum L Sp Pl 1099
(1753)

Mex & Ind. S Am

Trichomanes Preurii Kze.

251

1892-95 Common Rare in shady places along streams

Trichomanes L
T. elegans Rich. Act. Soc.
Hist Nat Paris 1. 114 (1792)
~~T. Brown~~ T Preurii Kze. Anal.
46 ~~(1738)~~ (1837)
W. Ind S. Am.

Trichomanes radicans Sw
Ceres

252
1526-28 On rotten tree trunks on shore
Nopu Bay

253
1465- On dead tree trunks at 600 ft

~~1528~~

254
1398 on side on wet perpendicular
cliff
Trichomanes L
T radicans. Sw. Schrad
Jour 1800, 2 97 (1801)
Widely distr. in both hemispheres

Acalypha bisetosa Spreng Euph
Cuba
Bon
13/4-15 Bushes 8 ft high on bank of
stream near shore

A. bisetosa Bert. acc. to
Spreng. Syst III. 879 (1826)

Porto Rico Venezuela.

Alsophila armata
Cocos

227
1340

Alsophila R. Br.

A. armata (Sw.) Pr. ~~Lunt~~
Lent. 62 (1836) Polypodium
armatum Sw. Prod. 134 (1788)

Jam. W Ind. Sam.

Asplenium ciciarium

228
1388-89

Common on wet rocks in a very restricted area side of a perpendicular cliff.

A cristatum Sam. Encycl II 310 (1786)

A Ceciarium Sw Prod 130 (1788

Mrs Hubbard Sam collected

Anona Cherimolia
Cocos
295
1573-76 In garden trees

Cornelius Mudflora
Cocos

~~1934~~
288
1332-34 On beach of stream near Chatham Bay.

Gymnogramme calomelanos Kaul.
Cocos.

230
1514 9/05

Cryptogramma Link,

C calomelanos (L.) Und. Bull.
Torr. Cl. XXIX 632 (1902)
Acrostichum calomelanos
L. Sp. Pl. 1072 (1753)
Gymnogramma calomelanos
Kf. Enum 76 (1824)
W Ind. Sam. Nat Afric

Lycopodium linifolium L Sp Pl!

Coca

255
1424 Specimen taken from 300 ft. Not as common as taken down

✓ 256
1476 Occasional at 600 ft

✓ 257
1293 On tree trunks

✓ 258
1342 on trunks of trees (This species do decrease in abundance as the

L Cinfolium L Sp Pl 1563 (1753)

Tree of Jud. N. Sam.

Nephrodium molle Sw?

Co—

1231
1361/64 9/11 specimen sterile & doubtful
1232
1472-73 9/11 Abundant 600 ft

Dryopteris adans.

D parasitica (L) O Ktze.
Rev. Gen II 811 (1891)
Polypodium parasiticum L
Sp Pl 1092 (1753)
Nephrodium molle Desv
Mem Soc. Linn V1 258 (1827)
Mex N Ind 8 am old Fld

Nephrolepis acuta
Coss

✓234
134 ~~4-43~~ 9/05 common in dense brakes 6-8°
high 250-300 ft Chatham Bay

✓235
1352-53 common in dense brakes 375-600 ft Chatham
Bay

✓236
1312 13 9/4 Roper Bay common in brake about
12 5 ft.

~~344~~

~~358~~

237
1316

Nephrolepis Schott,
N. biserrata (Sw) Schott,
Gen Fil ad. t 3 (1834
aspidium biserratum Sw Schrad.
Jour 1800 II, 32 (1801) Nephrolepis
acuta Presl, Tent. Pterid 79 (1836) Low

Mel Ward Sam Old
World

Nephrolepis pectinata
Schott.

238
1338

In vegetable mold side of
stream out

N. pectinata (Willd) Schott,
Gen Fil ad t 3 (1834)
aspidium pectinatum Willd
Sp Pl V 5 223 (1810)
Nephrolepis pectinata Schott,
Gen. Fil under t 3 (1834)

Phytolacca icosandra L Sp Pl 631

Orcas

V.J. 94
/309-11 occasional on banks of streams

Mex. N Ind. N S Am

Polypodium aureum
Coco

V-245-
1358-59 Common on sides of mountain
cliffs

P. aureum L. Sp. 1087 (1753)
widely distributed

Polypodium Phyllitides L Sp 1083

Coco

✓ 243
1365 - Common on trunks of trees

✓ 244
1367 " " " " "

Sub mus Hudson

Selaginella Galeottii Spring.

√ 258
1308

On banana trees at Nopeu
boy

Selaginella Blanc.
S Galeottii Spring. Monog
Lycopod 220 (1842-49)

Mex N. S. Am acc to
Spring l.c.

Setaria petoza
Cocu

264
1504.5. Common in cultivated ground
Nope Bay

Tell an doin sp
Close to J utriculata L but with
broader leaves and shorter flower
pedicels. a specimen (doubtful) of
labeled Cotopsis aloides B&K. in
herb. Gr. collected on Cocos Isl
by S & H & the Hopkins Stanford
defd probably in the 2nd

286
1363-64

287
1348 Common on trunks & branches of
trees

Anthurium Scandens Engl Dracontium

Cocos

277
1456 common on trees at 600 ft

Anthurium Schott.
A. scandens ~~Engl~~ (Aubl.) Engl.
~~(1915)~~ in Mart Fl Bras III. Pt 2
78 (1878-82) Dracontium scandens
Aubl. Pl. Gui. II. 836 (1775).

Cent. Am

Philodendron Schott, Aracea

280
1393 Occasional coming together and Small Tree near Stream Leading into Hopper Bay

~~Spathiphyllum~~ Wendlandii Schott. Araceae

~~(Engler)~~ Schott.

Cocos

√ 2 7 5
1278
2 7 6
√ 1293
~~1543~~
2 7 7
1300
√ 1541-43 2 7 8

Common on fork of stream occas 600 ft

S pathiphyllum Schott

S. Wendlandii Schott in Oesterr. Bot. Zeitschr VIII 179 (1858)

Cent. Amer.
~~Costa Rica~~

Sassafia Decne. Asclepiad.
J Colubrina Decne.

✓ 3 21
145·2 - 5-5 Abundant at 650 ft

(H.f. 7:12) Cuba Schomburgk

Sassafia Decne.
J Colubrina Decne. D C Prod VIII
5 79 (1844)

Brazil

Blainvillea biaristata DC Comp
Cucs

3 1 9
15-00-01 common in cultivated ground

B. biariotata DC Prod. V
#42(1836)

Brasil

Cliba dium accuminatum Benth.

Compositae

Coco

common bushes

326
1441

Cliba dium L.
C. accuminatum Benth. Bot.
Sulph. 114 (1844)

Endemic

Rolandra argentea Rottb (Comp
Cocos

327
1354

Rolandra Rottb
Rargentea Rottb. Coll. Hann.
11 2 58 (1775)

Panama W Ind N. Sam.

Acrocomia palmicola Cocos

328
1346-47
~~1347~~

Cocos
Abundant in open
places on hillsides
at Chatham Bay & Cocos
Gulf Nicaragua Bay

Acrocomia Jacq.

N. palmicola D.C. Prod V 538 (1836)

Nicaragua N.S. Am

Ipomoea Catharticа Poir. Convolvulaceae
Cocas

~322
1495-97 Common on open hillsides
covering bushes & other
vegetation with a heavy
~323
1544-51 growth

I catharticа Poir Diet
Supl. IV 633. (1816)

Florida. N. Ind. N. Sam.

Ipomea Pes-Caprae
Linn

324
1552-55- common on beach
~~1554~~ Hope Bay

Cyperus prolixus H B K

Cocos

268
1326

abundant on flat area near Playa
Agua Buena

Immature specimen doubtful as
to species. Seems to resemble
this one closely however

C prolixus H B K Nov. Gen.
Sp. 1. 206 (1815-)

Mex. N. Sam.

Cyperus sphacelatus Rottb.
Cocos

267
1327

288
1570-72 In low ground near Hofu Bay

~~1572~~

269
1506

C. sphacelatus Rottb. Descr.
& Ic. 26. (1786)

N. Ind N. S. Am.

Hymenophyllum Sp
Coed

233
1398-99 & 1405 Side 2 wet perpendicular
cliff above moss bay

Hymenophyllum Sm.

Hypolytrum Rich. n. sp. Cyperaceae

Cocos

V 2 7
1306

In Large bunches 5 ft or more
high Sec woodland & on bank
at Kaw Negeri Bay

271

1298-99 Abundant at 125 ft

Hypolytrum Rich.

Endemic.

Hypolytrum nicaraguense Liebm.

Chloris paniculata Schubler
& Cos from

✓ 260

1600-08 common on exposed cliffs along shore
& on the small islands

Chloris Sw.

C. paniculata Scribner,
in Rob. H. Gal. Isl. Proc
Am. Acad XXXVIII 704, 262
(1902)

Endemic

Panicum sanguinale
Coeds

2 Cr
V145-D-54
3262
1337

In cuvies of coeds along
stream leading into Hopi Bay
In open areas near Hopi Bay

Paspalum conjugatum. Grass

✓263
1355-

Common in open areas
around Roper Bay ✓

Clusia L rosea Jacq Enum. Pl Hyperoeae

Jacq. 34 (1760) Cocus

282

V/SGB-99

Common near Shoe

283

1324

Clusia L

C rosea ~~L Sp Pl 1145 (1753~~

Jacq Enum. 34 (1760)

Panama N Jud N Sam

Coesalpinia bonducella Leguminosae

297

1359 near shore Lumbini 6-Oct 1968

Cassia reticulata Willd Legum
Coes

1322 a clump of these bushes
occurs near the beach at
Chatham Bay

C. reticulata Willd. Enum Hort
Berol. 443 (1809)

Mex. N.S Amer.

Desmodium Sp Leguminosae
Cocos

298
1509-11 common at Rope Boy & Dampier Head

Acrostichum apodum. Filices

Cocos.

~~13201~~ 9/5- ~~Faicly~~
1321 9/5- On rotten logs not abundant.

Leptoglossum Schott,
E. apodum (Kl.) Scholly Gen
ad t. 14 (1834) Acrostichum
apodum Kl. Enum. 5·9 (1824)

W. Ind. Peru Brazil.

Elaphoglossum apodum

Acrostichum aureum Jacq.
2
S. Sp. Pl. 1069 (1753)
General in Trop.

Coco

225
1289. This fern grows to a height
of 6-8 ft sometimes as many
as 30 fronds in a bunch
Root stock 3 in in diameter
Fronds dense thickets in
places being the most
abundant undergrowth
extends up to 1285 ft in
wet woodland and seems
to leave off about where
1312 begins

Acrostichum cervinum Sw Fuloes

Cocas Island

1291-92 Sy4 Abundant in woodland on bank of stream leading into Hozen Bay. 2 to 3 ~~bogs~~ ft or more high

Polybotrya H & B

cervina (L.) Kl. Enum. 55 (1824). Osmunda cervina L Sp Pl 1065 (1753) acrostichum cervinum Sw Syn Fil. 14, 200 (1806)

S. Mex N Ind S. Am. to S Brazil

Adiantum Kaulfussii Kze. Filices
1632-35-8/12 Cocos.

" In crevices of rocks along
stream Wyed Bay

a petiolatum desv. Berl
mag V 326 (1811) a Kaulfussii
Kze. Linnaea XX 1 221 (1848)

Mex H and S Am

Aspidium Capense Willd? Filices
Cocos

241
323 9/4 Specimen is sterile & doubtful

242
1361

Polystichum Roth

Adiantiforme (Forst)
J. Sm. Hist Fil 220 (1875).

Polypodium adiantiforme
Forst Prod 82 (1786)

Asplenium coriaceum
Sw. Syn Fil 57 (1806)

Adiant. Sand. Del. wood

Olandra nodosa ~~Cav~~ Fil. ces.
Cocos

239
1480.81 9/8 Common on trees 600.

Olandra Cav.

O. nodosa (Willd.) Pr. Sent.
78 (1836) aspidium nodosum
Willd. Sp. Pl. v. 211 (1810).

cent Am. & Jud W. S. Am

Hibiscus tiliaceus Malvaceae
Cocos

301
1314

302
1351

303
1305.

304
1579-79

Clidemia hirta (L) D Don Melastom

Cocos

✓ 285-
1467-68 Common bushes & small trees at
600 ft

Clidemia D. Don.

C hirta (L.) D. Don. in Mem.
Wernerian Soc IV 309 (1823)
Melastoma hirta L Sp Pl 390 (1753)
Mex, Sam. (W Ind

Melastoma hirta Smith

Sw Obs

Dict no 3

175-

wild Sp. 2 5-68

Clidemia pustulata D.C. || Melastomaceae

1302-0 # Common bushes in woodland, about 8 ft high

C umbonata (Schr. & Mart) D.C. Prod. III. 158 (1828)
Melastoma umbonatum Schr & Mart. D C Prod l. c

Buzzk
Sam

Conostegia lasiopoda Benth Melastom.
Cocos

V306
1380 Small tree common

Conostegia D Don
C lasiopoda Benth Bot Sculph
9b (1444) Endemic

Miconia dodecandra Cogn Melastoma
Ocos

307
135-1

Miconia Ruiz + Pav.
~~Conostegia~~

M. dodecandra (Desv) Cogn in Mart
Fl. Bras XIV Pt 4 243 (1887)
Melastome dodecandra Desv. Jam
Ency. IV. 46 (1796)

Mex. N. Ind. N. S. Am.

Ossaea DC Macrophylla Cogn. Wilaston

308 Cocos
1474-75 a common tree at 600 ft

O. macrophylla Cogn. D.C
Monogr. VII 1064 (1891)

Endemic

Rapanea Guayanensis Aubl. Myrsinaceae

Coca

A 320
1402-03

Bushes about 8 ft high on bank of stream

~~Rapanea~~ a Rapanea Aubl.

R. Guianensis Aubl. Pl Gui 12(1775

Mex. Pl Ind. N. Sam.

Ardisia humilis Vahl Myrsinaceae
Cocos

V 313
1346-47

Knoches at 600 ft

A

Ardisia Sw.
a humilis Vahl.: Symb. III 40
(1794)
The specimen is fruiting so its exact
specific character is doubtfull resemb
a fruiting specimen of A humilis ih. Nth Fi, Cloudy
East Indies ~~Malay Archip~~

Eugenia Pacifica Beuth Myrtaceae
Cocos

1390-91 [2]84 Saw bushes common along streams & on hillsides

E. pacifica Beuth Bot Sulph. 98 (1844)

Endemic

Jussiaea linifolia Vahl. Onagrosa
Ceco

304
1356-57 Common covey tree & bushes

Jussiaea L.
J. linifolia Vahl. Ecolog.
Am 11 32 (1798)

Widely distr in trop

Palmar sp

1273
1461

1274
137 A

Peperomia nigra punctata Peperomia
Cocos

289
1384 Occasional on rotten
logs

P. nigra-punctata Miq Syst.
Pip. ~~##~~ 188(1840)
~~Dom~~

Martinica Isl.

Fleurya aestuans Gaud Urticaceae

Cocos

292

1395-96

common in open country

293

1414

round Moji Bay

Cornutia grandifolia Schauer (Verbenaceae)

368
133p-31 Fairly abundant on ridge
& cliffs & on banks of stream
Shrubs about 8 ft high

Cornutia L.
C. grandifolia ~~Ch & Schl.~~
~~Siphaea D~~ 97 (Ch & Schl.)
Schau. in DC Prod X 1682 (1847
Hosta grandifolia Ch & Schl.
Linn. V. 97 (1830)

S. Mex.